BETTER SAFE THAN SICK

TAKE PRECAUTIONS WHEN DEALING WITH BLOODBORNE & AIRBORNE PATHOGENS





- **1 FLUSH THE AREA** with water for 20 minutes
- 2 WASH EXPOSED AREAS using soap and water
- 3 TREAT SCABS & SORES GENTLY when cleaning your skin
- 4 REPORT THE EXPOSURE to your supervisor immediately
- 5 SAVE potentially contaminated objects so they can be tested
- 6 GET MEDICAL CARE as soon as possible

TO PROTECT AGAINST AIRBORNE PATHOGENS:

DROPLETS OF MOISTURE FROM AN INFECTED PERSON'S COUGH OR SNEEZE CAN REMAIN AIRBORNE FOR SEVERAL HOURS!

- ISOLATE PEOPLE with meningitis, influenza, pneumonia and tuberculosis
- 2 USE UV LIGHTING to destroy environmental bacteria
- 3 INSTALL SPECIAL AIR FILTERS where appropriate
- **WEAR** a fitted face mask and protective clothing

For more information on this and other safety topics, please log on to the member-exclusive website at **nsc.org**.

TOWAL OF A PROPERTY OF A PROPE

0216 900007681 ©2016 National Safety Council SOURCES: PDF 900006002 Proud Member